



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Combining relations (2) and (5) we obtain

$$k = q/4. \quad (6)$$

Equating the values of k given by relations (5) and (6) we find that p and q are connected by the linear equation

$$2p - q + 2 = 0. \quad (7)$$

Substituting k from relation (6) in relation (3) and then employing equation (7) we find

$$r = -3q^2/16 = -3(p+1)^2/4. \quad (8)$$

Incidentally, the last result shows that r must be negative in order that p and q may be real. Relations (7) and (8) enable the given quartic to be expressed in terms of (say) p alone, namely

$$x^4 + px^2 + 2(p+1)x - 3(p+1)^2/4 = 0,$$

or

$$[x^2 + (p+1)/2]^2 = [x - (p+1)]^2.$$

Consequently, the roots of the quartic are

$$x = (1 \pm \sqrt{-6p-5})/2,$$

$$x = (-1 \pm \sqrt{2p+3})/2.$$

For $-3/2 \leq p \leq -5/6$ all four roots will be real. For values of p outside of this range two roots will be real and two complex.

The reduction of the general quartic to the difference of two squares is given in Burnside and Panton's *Theory of Equations*, vol. I, p. 129, Art. 63 (4th or 7th editions.)

2783 [1919, 312]. Proposed by C. C. BRAMBLE, U. S. Naval Academy.

Two players, A and B , take turns throwing a single dice, A leading. The one first making a score of three aces is to be the winner. Find the probability that A will win.

SOLUTION BY H. P. MANNING, Brown University.

The probability of throwing three aces in the first three throws is $1/6^3$; in general, the probability of throwing exactly two aces in the first n throws, and an ace in the $(n+1)$ th throw, is $\frac{n(n-1)}{2} \cdot \frac{5^{n-2}}{6^{n+1}}$. That is, the probabilities of the different ways of throwing the first three aces are for A or B

$$\frac{1}{6^3}, \quad 3\frac{5}{6^4}, \quad 6\frac{5^2}{6^5}, \quad \dots$$

Call these numbers

$$a \quad b \quad c \quad \dots$$

Their sum should equal 1.

The probability that A wins is

$$a + (1-a)b + (1-a-b)c + \dots,$$

the factor in the parenthesis in any term indicating the probability that B has not yet thrown three aces. This expression may be written

$$a + b + c + \dots \\ - (ab + ac + bc + \dots).$$

Similarly the probability that B wins is

$$(1-a)a + (1-a-b)b + \dots,$$

which may be written

$$a + b + c + \dots \\ - (a^2 + ab + b^2 + ac + bc + c^2 + \dots).$$

Put

$$M = a + b + c + \dots$$

$$N = a^2 + b^2 + c^2 + \dots$$

$$P = ab + ac + bc + \dots,$$

$$M = \frac{1}{2 \cdot 6^3} \left[1 \cdot 2 + 2 \cdot 3 \cdot \frac{5}{6} + 3 \cdot 4 \left(\frac{5}{6} \right)^2 + \dots \right],$$

$$N = \frac{1}{4 \cdot 36^3} \left[(1 \cdot 2)^2 + (2 \cdot 3)^2 \frac{25}{36} + (3 \cdot 4)^2 \left(\frac{25}{36} \right)^2 + \dots \right].$$

Let

$$f(x) = 1 + x + x^2 + \dots = \frac{1}{1-x} \quad (|x| < 1);$$

then

$$f''(x) = 1 \cdot 2 + 2 \cdot 3x + 3 \cdot 4x^2 + \dots = \frac{2}{(1-x)^3}.$$

Hence putting $x = \frac{5}{6}$ we have $M = \frac{1}{2 \cdot 6^3} \cdot \frac{2}{(1/6)^3} = 1$.

$$x^2 f''(x) = 1 \cdot 2x^2 + 2 \cdot 3x^3 + 3 \cdot 4x^4 + \dots.$$

Differentiating twice,

$$2f'''(x) + 4xf''''(x) + x^2 f^{iv}(x) = (1 \cdot 2)^2 + (2 \cdot 3)^2 x + (3 \cdot 4)^2 x^2 + \dots.$$

This equals

$$2 \frac{2}{(1-x)^3} + 4x \frac{6}{(1-x)^4} + x^2 \frac{24}{(1-x)^5} = \frac{4}{(1-x)^5} (1 + 4x + x^2).$$

Hence putting $x = \frac{5}{6}$ we have

$$N = \frac{1}{4 \cdot 36^3} \cdot \frac{4 \cdot 36^5}{11^5} \left(1 + 4 \frac{25}{36} + \frac{25^2}{36^2} \right) = \frac{36^2 + 4 \cdot 25 \cdot 36 + 25^2}{11^5} = \frac{5521}{11^5}.$$

hence

$$M^2 = N + 2P = 1;$$

$$P = \frac{1}{2} - \frac{5521}{2 \cdot 11^5}.$$

Now the probability that A wins is $1 - P$ and the probability that B wins is $1 - N - P$. and so we may say, the probability that A wins is $N + P$ and the probability that B wins is P ; These probabilities are

$$\text{for } A \quad \frac{1}{2} + \frac{5521}{2 \cdot 11^5},$$

$$\text{" } B \quad \frac{1}{2} - \frac{5521}{2 \cdot 11^5}.$$

All of these series are absolutely convergent power series, the parentheses may be removed and the terms rearranged as shown in the expressions for the two probabilities, and M may be squared to give the relation $M^2 = N + 2P$.